

meducation[®]



0 1 5 T W U J 4



University Medical Center
123 Main Street
Anytown, USA 12345
Phone: (212) 555-1212

Foradil Aerolizer 12 mcg/inh

This medicine is used to treat asthma.

How to take medicine

Use the medicine each morning.

Inhale one (1) puff each time.



Morning	Noon	Evening	Bedtime
1 puff			

Instructions

The capsule needs to be placed into your inhaler before each use. Please follow the instructions for your inhaler carefully. Do not swallow the capsule.

Keep this medicine at room temperature. Protect from moisture and high humidity.

Ask your doctor, nurse or pharmacist to show you how to use this medicine correctly.

The medicine is a very fine powder. Some patients may not feel or taste the medicine. Do not take an extra dose of medicine if you do not taste or feel the medicine being delivered.

It is important that you keep taking each dose of this medicine on time even if you are feeling well.

If you forget to take a dose on time, take it as soon as you remember. If it is almost time for the next dose, do not take the missed dose. Return to your normal dosing schedule. Do not take 2 doses of this medicine at one time.

Please tell your doctor and pharmacist about all the medicines you take. Include both prescription and over-the-counter medicines. Also tell them about any vitamins, herbal medicines, or anything else you take for your health.

Do not share this medicine with anyone who has not been prescribed this medicine.

Cautions

Do not take the medicine more than twice during 24 hours.

Do not use the medication any more than instructed.

If this medicine causes more wheezing or makes it harder to breathe, stop the medicine. Get medical help right away.

Your ability to stay alert or to react quickly may be impaired by this medicine. Do not drive or operate machinery until you know how this medicine will affect you.

Tell the doctor or pharmacist if you are pregnant, planning to be pregnant, or breastfeeding.

Do not use this medicine during an asthma attack.

This medicine will not stop an asthma attack once it has started.

Do not start or stop any other medicines without first speaking to your doctor or pharmacist.

If you have any trouble using the medicine, please tell your doctor, nurse or pharmacist.

This medicine can cause serious side effects in some patients. Important information from the U.S. Food & Drug Administration (FDA) is available from your pharmacist. Please review it carefully with your pharmacist to understand the risks associated with this medicine.

Always keep medicine out of reach of children and pets.

Side Effects

- agitated feeling or trouble sleeping
- rarely, a severe episode of asthma may occur
- dizziness
- dry mouth
- lack of energy and tiredness
- increased risk for an infection
- nausea
- rapid heartbeat
- shakiness
- stomach upset or abdominal pain
- changes in taste or unpleasant taste

These are some potential side effects from this medicine. Please speak with your doctor about what you should do if you experience these or other side effects from your medication.

A few people may have an allergic reactions to this medicine. Symptoms can include difficulty breathing, skin rash, itching, swelling, or severe dizziness. If you notice any of these symptoms, seek medical help quickly.

Please speak with your doctor, nurse, or pharmacist if you have any questions about this medicine.

A demonstration is available online for this medicine.



IMPORTANT NOTE: This information is being provided to you as a supplement to other written instructions and package inserts provided by your doctor or pharmacist. This document contains a summary of your medication instructions and may not include all the product warnings or other important information about your medication. Please speak with your doctor or pharmacist about any question you may have about your medication or any unexpected symptoms. Always follow their advice and instructions.

